

Towards a diagrammatic derivation of the Veneziano-Yankielowicz-Taylor superpotential

J. Ambjørn^a and *R.A. Janik*^b

^a The Niels Bohr Institute, Copenhagen University
Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark.
email: `ambjorn@nbi.dk`

^c Institute of Physics, Jagellonian University,
Reymonta 4, PL 30-059 Krakow, Poland.
email: `ufrjanik@if.uj.edu.pl`

Abstract

We show how it is possible to integrate out chiral matter fields in $\mathcal{N} = 1$ supersymmetric theories and in this way derive in a simple diagrammatic way the $N_f S \log S - S \log \det X$ part of the Veneziano-Yankielowicz-Taylor superpotential.

Introduction

The recent renewed interest in the calculation of the glueball superpotential via matrix models [1] has led to an understanding of how to extract the non-logarithmic part of these superpotentials by ordinary diagrammatic methods [2]. Just as the matrix models in the applications to non-critical strings and 2d quantum gravity were convenient tools for solving specific combinatorial problems: the summation over all “triangulated” worldsheets with given weights, we understand now that the matrix model in the Dijkgraaf-Vafa (DV) context is an effective way of summing a set of ordinary Feynman graphs which by the magic of supersymmetry can be combined in such a way that they have no space-time dependence.

However, we are still left without a simple diagrammatic derivation of the logarithmic part of the glueball superpotential, the so called Veneziano-Yankielowicz-Taylor superpotential. This effective Lagrangian was originally derived for a pure $\mathcal{N} = 1$ $U(N_c)$ gauge theory by Veneziano and Yankielowicz [3] by anomaly matching and, by the same method, generalized to a $U(N_c)$ theory with N_f flavors in the fundamental representation by Taylor, Veneziano and Yankielowicz [4]. It is given by

$$W_{eff}^{VYT}(S, X) = W_{eff}^{VY}(S) + W_{eff}^{matter}(S, X) \quad (1)$$

where $W_{eff}^{VY}(S)$ is the pure gauge part

$$W_{eff}^{VY}(S) = -N_c S \log \frac{S}{\Lambda^3} \quad (2)$$

while $W_{eff}^{matter}(S, X)$ denotes the part coming from N_f flavors in the fundamental representation:

$$W_{eff}^{matter}(S, X) = N_f S \log \frac{S}{\Lambda^3} - S \log \frac{\det X}{\Lambda^2}. \quad (3)$$

In the above formulas S denotes the composite chiral superfield $\mathcal{W}_\alpha^2/32\pi^2$ and $X = \tilde{Q}Q$ is the $(N_f \times N_f)$ mesonic superfield, Q being the chiral matter field. In (2) and (3) Λ is an UV cut off. Usually this UV cut off is replaced by a renormalization group invariant scale Λ_M by use of the one-loop renormalization group:

$$\Lambda_M = \Lambda e^{-\frac{8\pi^2}{(3N_c - N_f)g^2}}. \quad (4)$$

The beautiful derivation of (1)-(3) by anomaly matching has always been somewhat antagonizing since a clear diagrammatic understanding is missing. It is summarized in the following citation from [5]: “Its [i.e. (1)-(4)] only *raison d’etre* is the explicit realization of the anomalous and non-anomalous symmetries of SUSY gluodynamics”.

In this letter we point out that there exists a simple diagrammatic derivation of (3). The derivation is inspired by diagrammatic techniques used in [2] and the observation that the DV-matrix models techniques could be extended to cover the case of superpotentials depending on mesonic superfields by considering the constrained (Wishart) matrix integrals [6]

$$\int DQD\tilde{Q} \delta(\tilde{Q}Q - X) = \frac{(2\pi)^{\frac{N(N+1)}{2}}}{\prod_{j=N-N_f+1}^N (j-1)!} (\det X)^{N-N_f} \quad (5)$$

and taking the large N limit.

Perturbative considerations

The matter contribution to the effective superpotential was shown in [2] to arise from the path integral

$$\int DQD\tilde{Q} e^{\int d^4x d^2\theta \left(-\frac{1}{2}\tilde{Q}(\square - i\mathcal{W}^\alpha \partial_\alpha)Q + W_{tree}(\tilde{Q}, Q) \right)} \quad (6)$$

where $\mathcal{W}^\alpha$ is an external field and $\partial_\alpha \equiv \frac{\partial}{\partial \theta^\alpha}$. If the quarks are massive ($W_{tree} = m\tilde{Q}Q$) then the above path integral reduces to a functional determinant which can be easily evaluated using the Schwinger representation:

$$\frac{1}{2} \int_{\frac{1}{\Lambda}}^{\infty} \frac{ds}{s} \int \frac{d^4p}{(2\pi)^4} \int d^2\pi_\alpha \exp(-s(p^2 + \mathcal{W}^\alpha \pi_\alpha + m)) \quad (7)$$

where we introduced an UV cut-off Λ . Due to fermionic integrations the result is

$$\frac{\mathcal{W}^2}{32\pi^2} \int_{\frac{1}{\Lambda}}^{\infty} \frac{ds}{s} e^{-ms} \quad (8)$$

which reduces for large Λ to

$$S \log\left(\frac{m}{\Lambda}\right) \quad (9)$$

At this stage one could integrate-in X to obtain (3). However, as “integrating-in” is in fact an assumption and we would like to obtain the desired result perturbatively, or more precisely: diagrammatically. To this end we impose the *superspace* constraint

$$X = \tilde{Q}Q \quad (10)$$

at the level of the path integral (6). This is done by introducing a Lagrange multiplier chiral superfield α . Since the antichiral sector does not influence the chiral superpotentials, we will perform a trick analogous to [2] and introduce an antichiral partner $\bar{\alpha}$ with a tree level potential $M\bar{\alpha}^2$. Thus we have

$$\int d^4x d^4\theta \bar{\alpha}\alpha + \int d^4x d^2\theta M\bar{\alpha}^2. \quad (11)$$

The path integral w.r.t. $\bar{\alpha}$ is Gaussian and yields (c.f. [2])

$$-\frac{1}{2M} \int d^4x d^2\theta \alpha \square \alpha. \quad (12)$$

The final path integral is

$$\int D\alpha D\tilde{Q}DQ e^{\int d^4x d^2\theta \left(-\frac{1}{2}\tilde{Q}(\square - i\mathcal{W}^\alpha \partial_\alpha)Q - \frac{1}{2}\alpha \square \alpha - \alpha X + \alpha \tilde{Q}Q\right)}, \quad (13)$$

where we also took $W_{tree} = 0$ and fixed the auxiliary mass $M = 1$ (it will be clear from the arguments below that the result is independent of M).

This is no longer a free field theory, but nevertheless there are significant simplifications if we only want to extract the $\text{tr } \mathcal{W}^2$ dependence. This implies that we must have two $\mathcal{W}$ insertions per $\tilde{Q}Q$ loop. The integrals over the fermionic momenta thus force all graphs which contain an α -line in a loop to vanish. Thus we are left with graphs coming from (13) which have the structure of $\tilde{Q}Q$ loops connected by at most one α propagator, and α propagators connected to the external field X as shown in fig. 1.

Moreover, if the field X contains a zero momentum component, which will generically be the case, the integrals will be dominated by this constant mode which forces the α propagators to be evaluated at zero momentum. Consequently we have to introduce an IR cut-off Λ_{IR} . Each 0-momentum α propagator will then just contribute a factor of $1/\Lambda_{IR}$. Thanks to the above property we may find the full $\tilde{Q}Q$ propagator in terms of the α 1-point function which we will denote by F :

$$\frac{1}{p^2 + \mathcal{W}^\alpha \pi_\alpha + F}, \quad (14)$$




Figure 1: Only tree level graphs survive, i.e. we are left with the the graphs shown in fig. 1c).

and the effective action will be given by the formula (7) with m substituted by F :

$$S \log \det \frac{F}{\Lambda} \quad (15)$$

It remains to determine F . The Schwinger-Dyson equation for F is (see fig. 2)

$$F = -\frac{1}{\Lambda_{IR}} X + \frac{1}{\Lambda_{IR}} \frac{S}{F} \quad (16)$$

where we used

$$\int_0^\infty ds \int \frac{d^4 p}{(2\pi)^4} \int d^2 \pi_\alpha e^{-s(p^2 + \mathcal{W}^\alpha \pi_\alpha + F)} = \frac{S}{F} \quad (17)$$

Eq. (16) is quadratic and has 2 solutions. Since the final result has to be IR finite, we will take the solution which has a finite limit as $\Lambda_{IR} \rightarrow 0$. Therefore

$$F = \frac{S}{X} \quad (18)$$

and by substituting this back in (15) one obtains the desired result:

$$S \log \det \frac{SX^{-1}}{\Lambda}, \quad (19)$$

or, in the case of N_f flavors:

$$N_f S \log \frac{S}{\Lambda^3} - S \log \det \frac{X}{\Lambda^2}. \quad (20)$$

$$--- \text{---} \bigcirc_{\text{shaded}} = --- \text{X} + --- \bigcirc_{\text{empty}}$$

Figure 2: The Schwinger-Dyson equation for F .

Further examples

Exactly the same technique can be adapted to the theories studied in [7] where the matter effective superpotentials in terms of only mesonic fields are quite complex (see eqn. (1.1) in [7]) and follow from quite intricate physical analysis. However, as noted in [7] the superpotentials with both glueball fields and matter fields are simpler. The pure matter superpotentials can then be obtained by integrating out the glueball fields S_i .

The simplest case considered in [7] is a gauge theory with gauge group $SU(2)_1 \times SU(2)_2$, with a bifundamental matter field Q in the (2,2) representation. The natural gauge invariant matter superfield is

$$X = Q^2 \equiv \frac{1}{2} Q_{ab} Q_{cd} \varepsilon^{ac} \varepsilon^{bd}, \quad (21)$$

and the matter part of the superpotential $W_{eff}(S, X)$ is (eq. (4.19) in [7]):

$$(S_1 + S_2) \log \frac{S_1 + S_2}{X \Lambda} \quad (22)$$

We will now show that the expression (22) also follows from a diagrammatic reasoning.

Since for $SU(2)$ the fundamental and antifundamental representations are equivalent through $\tilde{Q}_a \equiv Q_{a'} \varepsilon^{a'a}$ the Lagrangian for the bifundamental fields takes the form:

$$Q_{a'b'} \varepsilon^{a'a} \varepsilon^{b'b} (\square - i \mathcal{W}_{ac}^{(1)\alpha} \partial_\alpha - i \mathcal{W}_{bd}^{(2)\alpha} \partial_\alpha) Q_{cd} \quad (23)$$

Again we introduce a Lagrange multiplier superfield α enforcing the above constraint. We thus have

$$Q(\mathbf{C} \otimes \mathbf{C})(\square - \mathcal{W}^{(1)\alpha} \otimes \mathbf{1} \pi_\alpha - \mathbf{1} \otimes \mathcal{W}^{(2)\alpha} \pi_\alpha + \frac{1}{2} \alpha) Q - \alpha X \quad (24)$$

where $\mathbf{C}^{ab} \equiv \varepsilon^{ab}$.

The analogue of formula (15) will then be

$$\frac{1}{2} 2(S_1 + S_2) \log \left(\frac{F}{2\Lambda} \right) \quad (25)$$

where the $1/2$ comes from the fact that we are dealing with a real representation, while the 2 comes from performing the trace over the trivial factor in $(\mathcal{W}^{(1)} \otimes \mathbf{1})^2$. The Schwinger-Dyson equation for F will then have the form

$$F = -\frac{1}{\Lambda_{IR}} X + \frac{1}{\Lambda_{IR}} \frac{1}{2} \frac{2(S_1 + S_2)}{F/2} \quad (26)$$

hence

$$F = \frac{2(S_1 + S_2)}{X} \quad (27)$$

Inserting F into (25) reproduces precisely the nontrivial result (22).

Another example studied in [7] for the gauge group $SU(2)_1 \times SU(2)_2$ is matter $L\pm$ in the (1,2) representation. The classical D-flat direction is labeled by $Y = L_{\alpha+} L_{\beta-} \varepsilon^{\alpha\beta}$ and the matter contribution to W_{eff}^{VYT} was found in [7] to be:

$$S_2 \log \frac{S_2}{Y\Lambda}. \quad (28)$$

We can also reproduce this expression¹ by computing diagrammatically the contribution from the $L_{\pm}$ fields, starting with the Lagrangian

$$L(\mathbf{C} \otimes \mathbf{1})(\square - \mathcal{W}^{(2)\alpha} \otimes \mathbf{1}\pi_{\alpha} + \alpha\mathbf{1} \otimes \mathbf{C})L - \alpha Y, \quad (29)$$

where the second component in the tensor product is the flavor space.

Discussion

We have shown that it is possible to obtain the matter part of some generalized $W_{eff}^{VYT}(X, S)$ potentials by simple diagrammatic reasoning. It would be interesting to generalize the diagrammatic derivation to the gauge part of the Taylor-Veneziano-Yankielowicz superpotential. That would complete the diagrammatic derivation of the glueball superpotential.

¹Up to a trivial rescaling of Λ . Note that in our approach the definition of the UV cut-off Λ (see e.g. (8)) is a matter of convention and may be modified.

Acknowledgments JA and RJ acknowledge support by the EU network on “Discrete Random Geometry”, grant HPRN-CT-1999-00161, RJ was partially supported by KBN grant 2P03B09622 (2002-2004) and J.A. partially supported by “MaPhySto”, the Center of Mathematical Physics and Stochastics, financed by the National Danish Research Foundation.

References

- [1] R. Dijkgraaf and C. Vafa, *Matrix models, topological strings, and supersymmetric gauge theories* Nucl.Phys.B644:3-20,2002, [hep-th/0206255]; *On geometry and matrix models*, Nucl.Phys.B644:21-39,2002, [hep-th/0207106]; *A perturbative window into nonperturbative physics*, [hep-th/0208048].
- [2] R. Dijkgraaf, M.T. Grisaru, C.S. Lam, C. Vafa and D. Zanon, *Perturbative computation of glueball superpotentials*, [hep-th/0211017].
- [3] G. Veneziano and S. Yankielowicz, *An effective Lagrangian for the pure $N=1$ supersymmetric Yang-Mills theory*, Phys.Lett.113B:231-235,1982.
- [4] Taylor, G. Veneziano and S. Yankielowicz, *Supersymmetric QCD and its massless limit: an effective Lagrangian analysis*, Nucl.Phys.B218:493-505,1983.
- [5] M. A. Shifman and A. I. Vainshtein, *Instantons versus supersymmetry: fifteen years later*. ITEP Lectures in Particle Physics and Field Theory. Edited by M. Shifman. Singapore, World Scientific, 1999. Vol. 2, pp. 485-648, [hep-th/9902018]
- [6] Y. Demasure and R. A. Janik, *Effective matter superpotentials from wishart random matrices*, Phys.Lett.B553:105-108,2003, [hep-th/0211082].
- [7] K. A. Intriligator, R.G. Leigh and N. Seiberg, *Exact superpotentials in four-dimensions*, Phys.Rev.D50:1092-1104,1994, [hep-th/9403198]